Blood pressure and blood sugar — managing the two big numbers

Behtash Saeidi, MD

Diabetes and high blood pressure are two of the most common chronic conditions, they often travel together, and both are largely silent — people usually feel completely well while damage is building. The good news: both respond strongly to everyday habits, and the "numbers to aim for" are not one-size-fits-all — they shift with age and overall health.

PART 1 — BLOOD SUGAR (DIABETES)

What is really going on

- Food is broken down into glucose (sugar) in the blood. **Insulin** is the hormone that acts like a key, letting sugar into cells for energy.
- In **type 2 diabetes**, the body stops responding well to insulin (insulin resistance) and eventually cannot make enough. Sugar backs up in the blood.
- Persistently high sugar quietly damages small and large blood vessels — this is what harms eyes, kidneys, nerves, heart, and brain over years.
- Key point: the goal of treatment is not just a number — it is preventing that long-term vessel damage.

How we measure control

- **A1C** is a blood test that reflects the average blood sugar over about 3 months — think of it as the "90-day report card."
- Many people now use **continuous glucose monitors (CGM)**, where the goal is "time in range" (how much of the day sugar sits in the healthy zone)— generally **over 70%** of the day for most adults

Goals of therapy BY AGE / HEALTH STATUS

targets are individualized — tighter for the young and healthy, more relaxed for the older and frail, mainly to avoid dangerous low sugars (hypoglycemia).

- **Younger / healthy adults, long life expectancy:** A1C **under 7%**, and as low as **under 6.5%** if achievable safely
- **Most nonpregnant adults:** A1C **under 7%**
- **Healthy older adults (65+):** around **under 7–7.5%**.
- **Older adults with several illnesses or frailty:** more relaxed, about **under 8%**.
- **Very frail / limited life expectancy / advanced dementia:** no strict A1C number — the focus shifts to avoiding symptoms of high and low sugar and keeping quality of life
- Teaching point: an A1C that is too LOW in an older person can be dangerous (falls, confusion from low sugar). Looser is safer in frailty.

Lifestyle changes (the foundation for everyone)

- **Food:** No single "diabetes diet." Emphasize vegetables, fruit, whole grains, lean protein; limit sugary drinks and ultra-processed foods. A Mediterranean-style pattern is well supported
- **Activity:** At least **150 minutes per week** of moderate activity (e.g., brisk walking), plus **breaking up long sitting** — even a 5-minute movement break each hour helps
- **Weight:** Losing **at least 5%** of body weight improves sugar, blood pressure, and cholesterol; **10% or more**, especially early, can even push type 2 diabetes into remission
- **Sleep and smoking:** Adequate sleep and quitting smoking are part of the plan

When medication is added

- Lifestyle is started at diagnosis for everyone — medication is added when lifestyle alone does not reach the individual's target, and sometimes right away if sugar is very high.
- Reassure the audience: starting medication is not a "failure." Diabetes is progressive, and medicines (many of which also protect the heart and kidneys) are tools, not punishments.

PART 2 — BLOOD PRESSURE (HYPERTENSION)

What is really going on

- Blood pressure is the force of blood pushing against artery walls. The top number (systolic) is the heart beating; the bottom number (diastolic) is the heart resting.
- When pressure stays high, the heart works harder and arteries stiffen and get damaged — raising the risk of **stroke, heart attack, kidney failure, and even memory loss** over time.
- Key point: it is called the "silent killer" because there are usually no symptoms until damage is done — which is why checking matters.

Goals of therapy BY AGE

- **General target: under 130/80 mm Hg** for most adults, including healthy older adults, if tolerated
- **Older / frail (80+):** be more cautious, individualize, and watch for dizziness, falls, and lightheadedness on standing.
- Teaching point: like with sugar, in the very elderly or frail, pushing pressure too low can cause harm — the right number balances benefit against side effects.

Lifestyle changes (the cornerstone)

Ranked roughly by how effective they are:

- **DASH eating plan** — rich in fruits, vegetables, low-fat dairy, whole grains — the single most effective diet for lowering pressure
- **Cut salt (sodium):** aim under **2,300 mg/day**, ideally toward **1,500 mg/day**
- **More potassium** from fruits/vegetables
- **Weight loss:** roughly **1 mm Hg drop for every 1 kg (2.2 lb) lost**
- **Exercise:** regular aerobic activity (brisk walking most days) plus some strength training.
- **Alcohol:** limit to **≤1 drink/day (women), ≤2/day (men)**, less is better.
- **Stress reduction:** breathing techniques, meditation, yoga as helpful add-ons.

When medication is added

- Everyone starts with lifestyle. Medication is added when pressure stays above target despite those efforts, or started right away when pressure is high enough to pose immediate risk (e.g., 140/90 or above, especially with diabetes or other risk factors).
- Common, safe, well-studied first medicines include ACE inhibitors/ARBs, calcium-channel blockers, and diuretics (water pills); more than one low-dose pill is often better tolerated than one high-dose pill

PART 3 — WHY THEY MATTER TOGETHER

- Diabetes and high blood pressure multiply each other's risk to the heart, kidneys, and eyes — controlling both is far more powerful than controlling either alone.
- The same core habits — healthy diet, moving more, losing excess weight, less salt, no smoking — improve BOTH at once.
- Final takeaways for the audience:
    1. Know your numbers (A1C and blood pressure).
    2. Targets depend on your age and overall health — ask what YOURS should be.
    3. Lifestyle is the foundation; medication is a helpful tool, not a failure.
    4. Because both are silent, regular checks are what keep you safe.